

Chapter one

"Random Variables"

1- Th concept of Random variables

مفهوم المتغيرات العشوائية

Definitions:

Given a random experiment with sample space S , A random variable (r.v.) X is a function which assigns to each element $w \in S$ a real number $X(w)$ in the set E .

Ex) a _ Suppose that an experiment consists of tossing two fair coins :

SoL) $S = \{HH, HT, TH, TT\}$ ____ random experiment

Random variables

(H) : تمثل عدد مرات ظهور
 $X = 0, 1, 2$
 (T) : تمثل عدد مرات ظهور
 $Y = 0, 1, 2$

$P(X = x)$

$$P(X = 0) = \frac{1}{4}$$

$$P(X = 1) = \frac{2}{4}$$

$$P(X = 2) = \frac{1}{4}$$



$$\rightarrow \sum P(X = x_i) = 1$$

X Values

$HH \rightarrow 2$

$HT \rightarrow 1$

$TH \rightarrow 1$

$TT \rightarrow 0$

c _ Suppose that an experiment consists of tossing a pair of unbiased dice , X be a random variable indicating the sum of the two faces :-

Sol/

X =	2	3	4	5	6	7	8	9	10	11	12
	(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)	(2,6)	(3,6)	(4,6)	(5,6)	(6,6)
		(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(3,5)	(4,5)	(5,5)	(6,5)	
			(3,1)	(3,2)	(3,3)	(3,4)	(4,4)	(5,4)	(6,4)		
				(4,1)	(4,2)	(4,3)	(5,3)	(6,3)			
					(5,1)	(5,2)	(6,2)				
						(6,1)					

$$P(X=x) = \frac{1}{36}, \frac{2}{36}, \frac{3}{36}, \frac{4}{36}, \frac{5}{36}, \frac{6}{36}, \frac{5}{36}, \frac{4}{36}, \frac{3}{36}, \frac{2}{36}, \frac{1}{36}$$

X	2	3	4	5	6	7	8	9	10	11	12
P(X=x)	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

And $\sum P(X=x_i) = 1$

ويمكن تمثيل (r.v.) وفق دوال وعلى النحو الاتي :-

$$P(X) = \begin{cases} 0 & x=0 \\ \frac{x-1}{36} & x=2,3,4,5,6,7 \\ \frac{13-x}{36} & x=8,9,10,11,12 \\ 1 & \sum P(X) = 1 \end{cases}$$

ويمكن تمثيل دالة خطية عند رمي زار مرة واحدة

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$P(X) = \begin{cases} \frac{1}{6} & x = 1, 2, 3, \dots, 6 \\ 0 & \text{other wise} \end{cases}$$

Random Variable (r.v.)

متقطع
discrete (r.v.)

$$X = 0, 1, 2, 3, 4, \dots$$

Probability mass function

دالة كثافة احتمالية

(P.m.f.) مزلة

حكمة $P(X)$ في انبعاث



$$1- P(X) \geq 0 \quad \text{و} \quad \sum_{i=1,2,3,\dots}^n P(X_i) \leq 1$$

$$2- \sum_{i=1}^n P(X_i) = 1$$



$$F(x) = \sum_{X_i \leq x} P(X_i)$$

مستمر
Continuous (r.v.)


$$a < X < b$$

Probability density function

دالة كثافة احتمالية

(P.d.f.) مزلة

حكمة $f(x)$



$$f(x) \geq 0 \quad \text{و} \quad 0 \leq f(x) \leq 1$$

$$\int_{-\infty}^{\infty} f(x) dx = 1$$



$$F(x) = \int_{-\infty}^x f(x) dx$$

دالة الاحتمال التراكمية
Cumulative Probability function

2- Distribution Function (D.F.)

دالة التوزيع

Definitions:

If X is a random variable (r.v.) defined on the sample space S , the cumulative distribution function (c. d. f.) or cumulative probability function (c. p. f.) of a discrete random variable, denoted by $F(x)$ & is defined by :-

$$F(x) = P(X \leq x) \text{ for any real number}$$

$$= \sum P(x_i)$$

Properties :


The distribution function $F(x)$ has following properties :-

$$1_ F(\infty) = \lim_{x \rightarrow \infty} F(x) = 1$$

$$2_ F(-\infty) = \lim_{x \rightarrow -\infty} F(x) = 0$$


$$0 \leq F(x) \leq 1$$

$$3_ F(x_1) \leq F(x_2) \quad ; \quad \text{if } x_1 \leq x_2$$



$$4_ P(a < x \leq b) = F(b) - F(a) \quad \text{for all } a < b$$

$$P(x > b) = 1 - F(b)$$



Ex)

$$P(x) = \begin{cases} \frac{x}{6} & , \quad x = 1, 2, 3 \\ 0 & , \quad \text{other wise} \end{cases}$$

Find :

1_ Probability mass function

2_ cumulative probability function


4

Sol) 1-

$$\begin{array}{c|ccc} x & 1 & 2 & 3 \\ \hline P(X=x) & 1/6 & 2/6 & 3/6 \end{array} \quad \sum P(X=x_i) = 1$$

$$\sum P(X=x_i) = P(X=1) + P(X=2) + P(X=3) = \frac{1}{6} + \frac{2}{6} + \frac{3}{6} = \frac{6}{6} = 1$$

تحقق الشرط



رسم الدالة

$$2_ F(X) = P(X \leq x)$$

$$F(0) = P(X \leq 0) = P(X=0) = 0$$


$$F(1) = P(X \leq 1) = P(X=0) + P(X=1) = \frac{1}{6}$$

$$F(2) = P(X \leq 2) = P(X=0) + P(X=1) + P(X=2) = \frac{1}{6} + \frac{2}{6} = \frac{3}{6}$$

$$F(3) = P(X \leq 3) = P(X=0) + P(X=1) + P(X=2) + P(X=3)$$

$$= \frac{1}{6} + \frac{2}{6} + \frac{3}{6} = \frac{6}{6} = 1$$

$$\therefore F(X) = \begin{cases} 0 & x < 1 \\ \frac{1}{6} & 1 \leq x < 2 \\ \frac{3}{6} & 2 \leq x < 3 \\ 1 & x \geq 3 \end{cases}$$



رسم الدالة

5

وفي حالة توفر الدالة التراكمية فقط والمطلوب إيجاد دالة الكتلة الاحتمالية $P(X)$ يتم اجراء عملة الطرح من الأعلى الى الأسفل وفق الأسلوب الاتي :-

$$- P(0 < x \leq 1)$$

$$P(X=1) = F(b) - F(a) = \frac{1}{6} - 0 = \frac{1}{6} \quad P(a < x \leq b)$$

$$- P(1 < x \leq 2)$$

$$P(X=2) = F(2) - F(1) = \frac{3}{6} - \frac{1}{6} = \frac{2}{6}$$

$$- P(2 < x \leq 3)$$

$$P(X=3) = F(3) - F(2) = 1 - \frac{3}{6} = \frac{3}{6}$$